arXiv:1212.1534v1 [hep-ph] 7 Dec 2012

Axial-vector mesons from $\tau \to AP\nu_\tau$ decays

G. Calderón*
Facultad de Ingeniería Mecánica y Eléctrica, Universidad Autónoma de Coahuila, C.P. 27276, Torreón, Coahuila, México

J. H. Muñoz† and C. E. Vera‡
Departamento de Física, Universidad del Tolima A. A. 546, Ibagué, Colombia

Axial-vector mesons $a_1(1260)$, $f_1(1285)$, $h_1(1170)$, $K_1(1270)$ and $K_1(1400)$ can be produced in semileptonic $\tau \to AP\nu_\tau$ decays, where P stands for the pseudoscalar mesons π or K. We calculate the branching ratios based in a meson dominance model. The exclusive channels $\tau^- \to a_1(1260)^- \pi^0 \nu$, $\tau^- \to a_1(1260)^0 \pi^- \nu$, $\tau^- \to f_1(1285)\pi^- \nu$ and $\tau^- \to h_1(1170)\pi^- \nu$ turn out to be of order $\mathcal{O}(10^{-4})$ and channels $\tau^- \to K_1(1270)^- \pi^0 \nu$, $\tau^- \to K_1(1270)^0 \pi^- \nu$, $\tau^- \to K_1(1400)^- \pi^0 \nu$ and $\tau^- \to K_1(1400)^0 \pi^- \nu$ of order $\mathcal{O}(10^{-6})$. These results indicate that the branching ratios could be measured in experiments.

PACS numbers: 12.40Vv, 13.35Dx

---

* german.calderon@uadec.edu.mx
† jhmunoz@ut.edu.co
‡ cvera@ut.edu.co

I. INTRODUCTION

Recently the BABAR Collaboration [1] performed measurements of the branching fractions of three-prong and five-prong τ decay modes. Specifically, the branching fraction for the $\tau^- \to f_1(1285)\pi^-\nu_\tau$ decay is measured

$$\mathcal{B}(\tau^- \to f_1(1285)\pi^-\nu_\tau) = (3.79 \pm 0.25) \times 10^{-4}, \tag{1}$$

where the axial-vector meson $f_1(1285)$ is detected using the $f_1 \to 2\pi^+2\pi^-$ and $f_1 \to \pi^+\pi^-\eta$ modes. We have averaged in quadrature the two branching ratios and added the statistical and systematic errors reported by BABAR Collaboration [1]. This result supersedes a value reported by BABAR Collaboration [2]. Besides CLEO Collaboration [3] published a branching ratio of $(5.8^{+1.4}_{-1.3} \pm 1.8) \times 10^{-4}$, which is also based on the $f_1 \to 2\pi^+2\pi^-$ mode.

The measurements of $\tau \to AP\nu$ decays can give information about the hadronic matrix elements $\langle AP|J_\mu|0\rangle$ at the moderate energy regime of τ decays. This can be relevant in nonleptonic two-body D and B decays, where the annihilation contribution plays a relevant role in the branching ratios and CP asymmetries, see Refs. [4–7].

The $\tau \to AP\nu$ decays can contribute to the τ decay modes with four-pseudoscalar mesons in the final state. The exclusive $\tau^- \to f_1(1285)\pi^-\nu$ decay contributes as intermediate state in τ decay involving five-pseudoscalar mesons. The former decays can be related to $e^+e^- \to (4\pi)^0$. The relation between the $\tau^- \to (4\pi)^-\nu$ decays and the electromagnetic annihilation $e^+e^- \to (4\pi)^0$ process appears because the components of the same current are present in both cases, i.e., the current $\bar{d}\gamma_\mu u$ in τ decay and $1/\sqrt{2}(\bar{u}\gamma_\mu u - \bar{d}\gamma_\mu d)$ in the case of e^+e^- annihilation. Thus, these two processes are related by isospin symmetry.

On the other hand, at theoretical level, the production of p-wave mesons in semileptonic τ decays has not been widely studied in the literature. In effect, Ref. [8] and this paper are the only works that have considered the production of orbitally excited mesons in $\tau \to MP\nu_\tau$ channels, where M can be an axial-vector meson or a tensor meson. The exception has been Ref. [9], where the branching ratio for the channel $\tau^- \to f_1(1285)\pi^-\nu$ estimated is 2.91×10^{-4}.

In this paper we estimate the branching ratios of the $\tau \to AP\nu$ decays, where P is the pseudoscalar meson π or K and A is the axial-vector meson $a_1(1260)$, $f_1(1285)$, $h_1(1170)$, $K_1(1270)$ or $K_1(1400)$. We have considered the kinematically allowed, the G-parity conserved and first-class currents decays. We use a meson dominance model, where the resonances that contribute are coming from the vector mesons $\rho(770)$, $\rho(1450)$ and $K^*(982)$ and the axial-vector meson $a_1(1260)$. The weak decay and strong coupling constants are determined from experimental data. We use flavor $SU(3)$ symmetry where there are no data to estimate the constants.

In our estimations, we find that the branching ratios for the exclusive channels $\tau^- \to a_1(1260)^-\pi^0\nu$, $\tau^- \to a_1(1260)^0\pi^-\nu$, $\tau^- \to f_1(1285)\pi^-\nu$ and $\tau^- \to h_1(1170)\pi^-\nu$ are of order $\mathcal{O}(10^{-4})$. The channels $\tau \to K_1(1270)\pi\nu$ and $\tau \to K_1(1400)\pi\nu$ decays are of order $\mathcal{O}(10^{-6})$. Finally, the modes $\tau \to a_1(1260)K\nu$ and $\tau \to K_1(1270)K\nu$ are kinematically suppressed. Up to now, the $\tau^- \to f_1(1285)\pi^-\nu$ decay is the only mode that can be compared with experimental measurements.

The paper is organized as follow. In Sec. II, we describe the amplitude in the meson dominance model and give the expression to estimate the width decay. The calculation of the weak decay and strong coupling constants are given in Sec. III. In Sec. IV, we present the estimation of the branching ratios for the modes considered in this paper. And finally, we conclude in Sec. V.

II. AMPLITUDE AND WIDTH DECAY FOR SEMILEPTONIC $\tau \to AP\nu$ DECAYS

The decay amplitude for the $\tau(p_\tau) \to A(p_A)P(p_p)\nu_\tau(p_\nu)$ process, where A denotes the axial-vector meson $a_1(1260)$, $f_1(1285)$, $h_1(1170)$, $K_1(1270)$ or $K_1(1400)$, and P is the pseudoscalar meson π or K, can be writing by

$$\mathcal{M}(\tau \to AP\nu_\tau) = \frac{G_F}{\sqrt{2}} V_{ui}\bar{u}(p_\tau)\gamma^\mu(1-\gamma_5)u(p_\nu)\langle A(p_A)P(p)|J_\mu(0)|0\rangle \ , \tag{2}$$

where G_F is the Fermi constant, V_{ui} (i=d or s) is the corresponding element of the Cabibbo-Kobayashi-Maskawa matrix and $J_\mu(0)$ is the $(V-A)$ weak current.

The hadronic matrix element $\langle A(p_A,\varepsilon)P(p)|J_\mu|0\rangle$ is parametrized by [10]

$$\begin{aligned}\langle A(p_A,\varepsilon)P(p)|J_\mu|0\rangle &= l(t)\epsilon_\mu + (\epsilon.p)[c_+(t)(p-p_A)_\mu + c_-(t)(p+p_A)_\mu] \\ &\quad + iq(t)\epsilon_{\mu\nu\rho\sigma}\epsilon^\nu(p_A-p)^\rho(p_A+p)^\sigma \ ,\end{aligned} \tag{3}$$

where the hadronic matrix element is expressed in terms of the form factors $l(t)$, $c_+(t)$, $c_-(t)$ and $q(t)$, which are Lorentz-invariant functions of squared momentum transfer $t = (p_A + p)^2$, the polarization vector ϵ_μ describes the axial-vector meson A.

To estimate the form factors, we use the meson dominance model, i.e., the decay amplitude is given by the sum of intermediate meson resonance contributions

$$\mathcal{M}(\tau \to AP\nu_\tau) = \sum_{R^*} \mathcal{M}(\tau \to R^* \ \nu_\tau \to AP\nu_\tau) \ , \tag{4}$$

where R^* is the intermediate resonance, which has the right quantum numbers for the specific processes. The sum is extended over all possible resonance contributions, see Fig. 1.

The production of the resonance R^* can be classified as either first- or second-class current depending on the spin J, parity P, and G-parity of the resonance particle. In the Standard Model, first-class current is considered to dominate. The second-class currents are associated with a decay constant proportional to the mass difference between an up and down quark and in the exact isospin limit symmetry they vanish [11]. The vector meson resonances $\rho(770)$, $\rho(1450)$, $K^*(982)$ and the axial-vector meson $a_1(1260)$ are produced by first-class currents, which are the contributions included in this work. The vertex $R^* \to AP$ decay, which is produced by strong interaction, must conserve G-parity.

Comparing the hadronic current, see Eq. (3), with the amplitude build from the meson dominance model, which is calculated using Feynman rules, we determine expressions for the form factors $q(t) = 0$ and $c_-(t) = -c_+(t) = c(t)$. Thus, the amplitude is expressed in terms of only two form factors $l(t)$ and $c(t)$. These form factors can be expressed by means of the Breit-Wigner (BW) function

$$\begin{aligned} l(t) &= -\frac{if_V g_{AVP}}{M} \ (p_A \cdot p_V) \ BW_V(t), \\ c(t) &= \frac{if_V g_{AVP}}{2M} \ BW_V(t), \end{aligned} \tag{5}$$

where $BW_V(t)$ is defined by

$$BW_V(t) = \frac{M^2}{M^2 - t - iM\Gamma} \ , \tag{6}$$

the parameters M and Γ are the mass and width decay of the intermediate meson resonant particle, where we have used the complex-mass prescription $M_0^2 \to M^2 - iM\Gamma$ and a constant width Γ.

The unpolarized squared amplitude in terms of the form factors $l(t)$ and $c(t)$ is the following

$$\sum_{pol} |\bar{\mathcal{M}}|^2 = 2G_F^2 |V_{ui}|^2 \Big[a_1(t,s)|l(t)|^2 + a_2(t,s)|c(t)|^2 + a_3(t,s) Re[l^*(t)c(t)] \Big], \tag{7}$$

where the kinematic factors are defined by

$$\begin{aligned} a_1(t,s) &= \frac{1}{m_A^2} \Big[(s - m_\tau^2)(s + t - m_\tau^2 - m^2) + m_A^2(m^2 + 2m_\tau^2 - 2t - s) \Big], \\ a_2(t,s) &= \frac{1}{m_A^2} \Big[((m^2 - t)^2 - 2(m^2 + t)m_A^2 + m_A^4)((m^2 - s)m_A^2 + (s - m_\tau^2)(s + t - m_\tau^2 - m^2)) \Big], \\ a_3(t,s) &= \frac{2}{m_A^2} \Big[(m^2 - s)m_A^4 + (m^2 - t)(s - m_\tau^2)(m^2 + m_\tau^2 - s - t) \\ &\quad + m_A^2(s^2 + m^2 t + m_\tau^2 t - m^4 - m^2 m_\tau^2 - m_\tau^4) \Big]. \end{aligned} \tag{8}$$

The kinematics factors $a_1(t,s)$, $a_2(t,s)$ and $a_3(t,s,)$ are functions of squared moments $t = (p_A - p)^2$ and $s = (p_\nu + p_A)^2$, and the masses of τ lepton m_τ , axial-vector meson m_A and pseudo-scalar meson m.

The width decay for the channel $\tau \to AP\nu_\tau$ is calculated integrating the unpolarized amplitude, Eq. (7), with respect to the variables t and s, and the appropriate kinematic limits.

The vector meson resonant contribution is the only contribution for the channels studies in this work, with the exception of the $\tau^- \to f_1(1285)\pi^-\nu$ decay, where it has an axial-vector meson $a_1(1260)$ as the only present contribution.



FIG. 1. Intermediate meson dominance resonance of $\tau \to AP\nu$ decays.

Specifically, for the channels $\tau \to a_1(1260)\pi\nu$, $\tau \to h_1(1170)\pi\nu$, $\tau \to K_1(1270)K\nu$ and $\tau \to K_1(1400)K\nu$ we have the contributions of vector meson $\rho(770)$ and $\rho(1450)$ resonances. The strange vector meson $K^*(982)$ contributes to the processes $\tau \to a_1(1260)K\nu$, $\tau \to K_1(1270)\pi\nu$ and $\tau \to K_1(1400)\pi\nu$.

In studies of the $\tau^- \to \pi^-\pi^0\nu$ channel carried out by the Belle [12] and ALEPH [13] Collaborations, was established that in order to describe correctly the spectral function it is necessary to introduce two resonances, the vector mesons $\rho(770)$ and $\rho(1450)$. We model the two contributions with a linear combination normalized by

$$BW_V(t) = \frac{BW_\rho(t) + \beta BW_{\rho'}(t)}{1+\beta}, \tag{9}$$

where the parameter β is determined by a fit to the hadronic spectral function obtained by the experiments. In our work, we model in the same manner the contribution of the two resonances $\rho(770)$ and $\rho(1450)$ to the channels considered above. We use a conservative value of $\beta = \pm(0.2 \pm 0.1)$, see Ref. [8].

III. WEAK DECAY AND STRONG COUPLING CONSTANTS

In order to obtain the weak decay and strong coupling constants that are required in this work, we make use of the available experimental information. However, for some channels, we use flavor $SU(3)$ and isospin $SU(2)$ symmetries to relate them to the constants obtained by experimental data.

We use the experimental reported branching ratios in Ref. [14] to determine the strong coupling constants from the decays of strange axial-vector mesons $K_1(1270) \to K\rho$, $K_1(1270) \to K^*(982)\pi$ and $K_1(1400) \to K^*(982)\pi$ and from the $h_1(1170) \to \rho\pi$ decay.

The strong coupling constant g_{AVP} (in GeV^{-1}) is determined from the amplitude of probability for the $A \to VP$ process, which is defined by

$$\mathcal{M}(A(p_A, \epsilon_A) \to V(p_V, \epsilon_V)P(p)) = g_{AVP}\epsilon_A^\mu \epsilon_V^\nu [(p_A.p_V)g_{\mu\nu} - (p_A)_\mu (p_V)_\nu] \ . \tag{10}$$

The decay rate in the rest frame of the decaying axial-vector meson is

$$\Gamma(A \to VP) = \frac{1}{3}\frac{g_{AVP}^2}{32\pi}\frac{\lambda^{1/2}(m_A^2, m_V^2, m^2)}{m_A^3}\left[2m_A^2 m_V^2 + (m_A^2 + m_V^2 - m^2)^2\right] \ , \tag{11}$$

where $\lambda(x,y,z)$ is the Kallen function, which is defined by $\lambda(x,y,z) = x^2 + y^2 + z^2 - 2(xy + xz + yz)$.

Using the measured width decay for the channels $K_1(1270) \to K\rho$, $K_1(1270) \to K^*\pi$, $K_1(1400) \to K\rho$, $K_1(1400) \to K^*\pi$, and using the isospin $SU(2)$ symmetry to relate with the charged modes, we obtain the following values in (GeV^{-1}): $g_{K_1(1270)^- K^0 \rho^-} = 5.04 \pm 0.67$, $g_{K_1(1270)^0 K^- \rho^-} = 5.04 \pm 0.67$, $g_{K_1(1270)^- K^{*-} \pi^0} = 0.69 \pm 0.13$, $g_{K_1(1270)^0 K^{*-} \pi^-} = 0.97 \pm 0.19$, $g_{K_1(1400)^- K^0 \rho^-} = 0.68 \pm 0.34$, $g_{K_1(1400)^0 K^- \rho^-} = 0.68 \pm 0.34$, $g_{K_1(1400)^- K^{*-} \pi^0} = 1.95 \pm 0.10$ and $g_{K_1(1400)^0 K^{*-} \pi^-} = 2.75 \pm 0.14$. The error reported is due to the experimental uncertainty in the width decay.

In order to extract the strong decay constant $g_{h_1 \rho\pi}$, we have assumed that only the mode $h_1(1170) \to \rho\pi$ contributes to the total width decay of the axial-vector meson $h_1(1170)$. We obtain $g_{h_1 \rho^- \pi^+} = 3.85 \pm 0.21$ GeV^{-1}. The uncertainty is determined by the experimental error in the width decay.

There are two types of axial-vector mesons, which in spectroscopy notation ${}^{2s+1}L_J$ are p-wave mesons ${}^3P_1$ and ${}^1P_1$. Considering the flavor $SU(3)$ symmetry, the ${}^3P_1$-nonet is integrated by $a_1(1260)$, $f_1(1285)$, $f_1(1420)$ and K_{1A} and the ${}^1P_1$-nonet composed by $b_1(1235)$, $h_1(1170)$, $h_1(1380)$ and K_{1B}. Here, in this work is relevant consider that the physical strange axial-vector mesons $K_1(1270)$ and $K_1(1400)$ are the result of a mixing of the flavor $SU(3)$ eigenstates K_{1A} and K_{1B} [15]. Thus, the mixing can be described by

$$\begin{aligned} K_1(1270) &= K_{1A}\sin\theta + K_{1B}\cos\theta \\ K_1(1400) &= K_{1A}\cos\theta - K_{1B}\sin\theta \ , \end{aligned} \tag{12}$$

where θ is the mixing angle. Restricting its value to $0 < \theta < 90°$, it is found two solutions $\theta = 32°$ and $\theta = 58°$, see Ref. [15].

We consider flavor $SU(3)$ symmetry to relate the strong coupling constants $g_{a_1\rho\pi}$ and $g_{a_1K^*K}$ with $g_{K_{1A}\rho K}$ and $g_{K_{1A}K^*\pi}$, respectively. The relations are $g_{a_1\rho\pi} = \sqrt{4/3}g_{K_{1A}K^*\pi}$ and $g_{a_1K^*K} = \sqrt{1/3}g_{K_{1A}\rho K}$.

Furthermore, we use isospin $SU(2)$ symmetry to relate the strong coupling constants for different charged states in the respective channels. For example, in the case of the axial-vector meson $K_1(1270)$, the branching ratio for the modes $K_1(1270) \to K^*\pi$ and $K_1(1270) \to K\rho$ are reported in Ref. [14]. The isospin relations are $g_{K_1^-K^0\rho^-} = \sqrt{2/3}g_{K_1K\rho}$, $g_{K_1^0K^-\rho^-} = \sqrt{2/3}g_{K_1K\rho}$, $g_{K_1^-K^{*-}\pi^0} = 1/\sqrt{3}g_{K_1K^*\pi}$ and $g_{K_1^0K^{*-}\pi^-} = \sqrt{2/3}g_{K_1K^*\pi}$. Similar analysis for strong coupling constant in decay modes for the axial-vector meson $K_1(1400)$ is done. The results obtained are: $g_{a_1^-\rho^-\pi^0} = 3.75 \pm 0.74$, $g_{a_1^0\rho^-\pi^-} = 3.75 \pm 0.74$, $g_{a_1^-K^{*-}K^0} = 1.17 \pm 0.09$, and $g_{a_1^0K^{*-}K^-} = 1.64 \pm 0.13$, in unit GeV^{-1}. The reported error is coming from the experimental uncertainty in the width decays.

Moreover we use the flavor $SU(3)$ symmetry to obtain the relation $g_{f_1a_1^-\pi^+} = 1/\sqrt{2}g_{a_1^0\rho^-\pi^-}$. Thus, the strong coupling constant has the value $g_{f_1a_1^-\pi^+} = 2.65 \pm 0.53\ \mathrm{GeV}^{-1}$.

We have estimated the weak decay constant of hadron mesons h^- from the experimental data on branching ratio from $\tau^- \to h^-\nu_\tau$ decays. We obtain $f_\rho = 218 \pm 2$ and $f_{K^*} = 210 \pm 5$ MeV. Also, weak decay constant for the physical states $K_1(1270)$ and $K_1(1400)$ are obtained. Using a mixing angle $\theta = 32°$ and inverting Eq. (12), we obtain the weak decay constant for the flavor K_{1A} and K_{1B} states. The weak decay constant for the axial-vector meson $a_1(1260)$ is $f_{a_1} = 213 \pm 10$ MeV, which is obtained by flavor $SU(3)$ symmetry from the weak decay of the mesons K_{1A} and K_{1B}, see Ref. [15]. In the calculations above, the error in the weak decay constants are coming from the experimental error in branching ratios of $\tau^- \to h^-\nu_\tau$ decays.

The necessary values for the relevant parameters such as width decays and masses of the particles involved in the channels calculated in this work have been taken from Ref. [14].

IV. BRANCHING RATIOS

The branching ratios for the exclusive $\tau^- \to AP\nu_\tau$ decays estimated in this work are shown in Table I.

The largest branching ratios estimated in this work turn out to be of order $\mathcal{O}(10^{-4})$, corresponding to the exclusive modes $\tau^- \to a_1(1260)^-\pi^0\nu$, $\tau^- \to a_1(1260)^0\pi^-\nu$, $\tau^- \to f_1(1285)\pi^-\nu$ and $\tau^- \to h_1(1170)\pi^-\nu$. In these modes we have the contribution of two vector meson resonances $\rho(770)$ and $\rho(1450)$, which produce a broad form factors $l(t)$ and $c(t)$ as function of the momentum transfered t, in consequence a large branching ratio estimation.

Up to now, the only measured branching ratio from the calculated in this work is for the channel $\tau^- \to f_1(1285)\pi^-\nu$. Our estimate for this channel is $(11.8 \pm 4.5) \times 10^{-4}$. We can compare with the recent result of BABAR Collaboration $(3.79 \pm 0.25) \times 10^{-4}$ [1] and determine that our estimation differs by approximately one sigma error with respect to the experimental result.

The channels with branching ratios of order $\mathcal{O}(10^{-6})$ are $\tau \to K_1(1270)\pi\nu$ and $\tau \to K_1(1400)\pi\nu$ decays. They have only the contribution of the vector meson $K^*(982)$ as intermediate resonance.

The exclusive channels $\tau^- \to K_1(1400)^0K^-\nu$ and $\tau^- \to K_1(1400)^-K^0\nu$ decays are not kinematically allowed. The modes $\tau \to a_1(1260)K\nu$ and $\tau \to K_1(1270)K\nu$ decays are of the order of $\mathcal{O}(10^{-9})$, this is due to the small phase space available for their decay, i.e, they are kinematically suppressed decays.

The channels $\tau \to a_1(1260)\pi\nu$, $\tau^- \to h_1(1170)\pi^-\nu$, $\tau \to K_1(1270)\pi\nu$ and $\tau \to K_1(1400)\pi\nu$ decays, with resonance contributions coming from the vector mesons $\rho(770)$ and $\rho(1450)$, have the largest error due to the error in the parameter β. In the other cases, where the meson resonance contribution is the strange vector meson $K^*(982)$ or the axial-vector meson $a_1(1260)$, the principal error in the branching ratio estimate is due to the uncertainty in the strong coupling constant g_{AVP}, where V is the resonance contribution $K^*(982)$ or $a_1(1260)$.

In addition, there are two other sources of error in our estimations not considered. We use flavor $SU(3)$ and isospin $SU(2)$ symmetries to relate the weak decay and strong coupling constants to the obtained by experimental information. The error in flavor $SU(3)$ symmetry can reach 20% in the estimation. The other source of theoretical uncertainty is coming from the contact term in the weak vertex of the hadronic current $\langle AP|J_\mu(0)|0\rangle$. However, we do not consider this contribution because it is associated with the continuum, i.e. there is no $BW_V(t)$. The contact term contribution is not possible to estimate in a meson dominance model.

TABLE I. Branching ratios for $\tau \to AP\nu$ decays.

AP mode	Branching ratio	Parameter β
$a_1(1260)^-\pi^0$	$(20.3 \pm 11.7) \times 10^{-4}$	$\beta = +0.2 \pm 0.1$
$a_1(1260)^-\pi^0$	$(25.4 \pm 18.8) \times 10^{-4}$	$\beta = -0.2 \pm 0.1$
$a_1(1260)^0\pi^-$	$(19.8 \pm 11.4) \times 10^{-4}$	$\beta = +0.2 \pm 0.1$
$a_1(1260)^0\pi^-$	$(24.5 \pm 18.2) \times 10^{-4}$	$\beta = -0.2 \pm 0.1$
$a_1(1260)^-K^0$	$(1.6 \pm 0.3) \times 10^{-9}$	
$a_1(1260)^0K^-$	$(4.6 \pm 0.6) \times 10^{-9}$	
$f_1(1285)\pi^-$	$(11.8 \pm 4.5) \times 10^{-4}$	
$h_1(1170)\pi^-$	$(29.1 \pm 16.6) \times 10^{-4}$	$\beta = +0.2 \pm 0.1$
$h_1(1170)\pi^-$	$(43.3 \pm 30.8) \times 10^{-4}$	$\beta = -0.2 \pm 0.1$
$K_1(1270)^-\pi^0$	$(8.3 \pm 3.0) \times 10^{-7}$	
$K_1(1270)^0\pi^-$	$(1.6 \pm 0.6) \times 10^{-6}$	
$K_1(1400)^-\pi^0$	$(1.3 \pm 0.2) \times 10^{-6}$	
$K_1(1400)^0\pi^-$	$(2.4 \pm 0.3) \times 10^{-6}$	
$K_1(1270)^0K^-$	$(11.6 \pm 5.8) \times 10^{-9}$	$\beta = +0.2 \pm 0.1$
$K_1(1270)^-K^0$	$(2.5 \pm 1.3) \times 10^{-9}$	$\beta = +0.2 \pm 0.1$

V. CONCLUSIONS

We have estimated branching ratios of the $\tau \to AP\nu$ decays. The axial-vector mesons A considered are $a_1(1260)$, $f_1(1285)$, $h_1(1170)$, $K_1(1270)$ and $K_1(1400)$. The pseudoscalar mesons P are π and K. In total, 12 exclusive channels are kinematically allowed, first class-currents kind and G-parity conserved.

To calculate the branching ratios we use a meson dominance model, where the form factors of the current $\langle AP|J_\mu|0\rangle$ are calculated from the vector meson contributions $\rho(770)$, $\rho(1450)$ and $K^*(982)$. With the exception of the channel $\tau^- \to f_1(1285)\pi^-\nu$, where the contribution is due to the axial-vector meson $a_1(1260)$.

The channels $\tau^- \to a_1(1260)^-\pi^0\nu$, $\tau^- \to a_1(1260)^0\pi^-\nu$, $\tau^- \to f_1(1285)\pi^-\nu$ and $\tau^- \to h_1(1170)\pi^-\nu$ decays are of order $\mathcal{O}(10^{-4})$. We can compare the estimated branching ratio $Br(\tau^- \to f_1(1285)\pi^-\nu) = (11.8 \pm 4.5) \times 10^{-4}$ with the measured value $(3.79 \pm 0.25) \times 10^{-4}$, recently reported by BABAR Collaboration [1].

We observe a pattern in the branching ratios calculated in this work. The modes of order $\mathcal{O}(10^{-9})$ are kinematically suppressed. The channels of order $\mathcal{O}(10^{-6})$ are Cabibbo suppressed and they have only the contribution of the vector meson $K^*(982)$. Finally, the channels with branching ratios of order $\mathcal{O}(10^{-4})$ are Cabibbo allowed and have two vector meson resonance contributions $\rho(770)$ and $\rho(1450)$, with the exception of the $\tau^- \to f_1(1285)\pi^-\nu$ where the resonance is the axial-vector meson $a_1(1260)$.

The processes studied in this work can contribute as intermediate states in branching ratios with four and five pseudoscalar mesons π and/or K in final states for the τ decay modes.

Eventually, the branching ratios for the channels calculated in this work, with the exception of the channel $\tau^- \to f_1(1285)\pi^-\nu$ (which have been already measured), will be measured by four-prong of τ decays, with the data sample of τ lepton pairs accumulated by the B-factories BABAR and Belle experiments [16]. Additionally, SuperB [17] and Belle-II [18, 19] Collaborations have programs to produce τ pairs and measure rare τ decays.

ACKNOWLEDGMENTS

G. C. acknowledges financial support from Promep, México. J. H. M. and C. E. V. are grateful to *Comité Central de Investigaciones* (CCI) of the University of Tolima, Colombia.

---

[1] J. P. Lees *et al.* [BABAR Collaboration], arXiv:1209.2734 [hep-ex].

[2] B. Aubert *et al.* [BABAR Collaboration], Phys. Rev. D **72**, 072001 (2005) [hep-ex/0505004].
[3] T. Bergfeld *et al.* [CLEO Collaboration], Phys. Rev. Lett. **79**, 2406 (1997) [hep-ex/9706020].
[4] H. -n. Li, C. -D. Lu and F. -S. Yu, Phys. Rev. D **86**, 036012 (2012) [arXiv:1203.3120 [hep-ph]].
[5] Z. -J. Xiao, W. -F. Wang and Y. -y. Fan, Phys. Rev. D **85**, 094003 (2012) [arXiv:1111.6264 [hep-ph]].
[6] Q. Chang, X. -W. Cui, L. Han and Y. -D. Yang, Phys. Rev. D **86**, 054016 (2012) [arXiv:1205.4325 [hep-ph]].
[7] X. Liu and Z. -J. Xiao, Phys. Rev. D **86**, 074016 (2012) [arXiv:1203.6135 [hep-ph]].
[8] G. L. Castro and J. H. Munoz, Phys. Rev. D **83**, 094016 (2011) [arXiv:1103.2993 [hep-ph]].
[9] B. A. Li, Phys. Rev. D **55**, 1436 (1997) [hep-ph/9606402].
[10] N. Isgur, D. Scora, B. Grinstein and M. B. Wise, Phys. Rev. D **39**, 799 (1989); D. Scora and N. Isgur, Phys. Rev. D **52**, 2783 (1995).
[11] S. Weinberg, Phys. Rev. **112**, 1375 (1958).
[12] M. Fujikawa *et al.* [Belle Collaboration], Phys. Rev. D **78**, 072006 (2008) [arXiv:0805.3773 [hep-ex]].
[13] S. Schael *et al.* [ALEPH Collaboration], Phys. Rept. **421**, 191 (2005) [hep-ex/0506072].
[14] J. Beringer *et al.* (Particle Data Group), Phys. Rev. D **86**, 010001 (2012).
[15] G. Nardulli and T. N. Pham, Phys. Lett. B **623**, 65 (2005).
[16] A. Lusiani, PoS HQL **2010**, 054 (2010) [arXiv:1012.3733 [hep-ex]].
[17] M. Bona *et al.* [SuperB Collaboration], Pisa, Italy: INFN (2007) 453 p. www.pi.infn.it/SuperB/?q=CDR [arXiv:0709.0451 [hep-ex]].
[18] T. Aushev *et al.* [BELLE II Collaboration], arXiv:1002.5012 [hep-ex].
[19] C. Z. Yuan [BELLE II Collaboration], arXiv:1208.3813 [physics.ins-det].